arXiv:1603.02072v1 [math.CA] 26 Feb 2016

ON PALEY-TYPE AND HAUSDORFF – YOUNG – PALEY-TYPE INEQUALITIES FOR GENERALIZED GEGENBAUER EXPANSIONS

Roman A. Veprintsev

Abstract. We establish Paley-type and Hausdorff – Young – Paley-type inequalities for generalized Gegenbauer expansions.

Key words and phrases: orthogonal polynomials, generalized Gegenbauer polynomials, Paley-type inequality, Hausdorff – Young – Paley-type inequality

MSC 2010: 33C45, 41A17, 42C10

1. Introduction and preliminaries

In this section, we introduce some classes of orthogonal polynomials on $[-1,1]$, including the so-called generalized Gegenbauer polynomials. For a background and more details on the orthogonal polynomials, the reader is referred to [2–4, 7].

Let $\mathbb{N}_0$ denote the set of non-negative integers. Let α, $\beta > -1$. The Jacobi polynomials, denoted by $P_n^{(\alpha,\beta)}(\cdot)$, $n \in \mathbb{N}_0$, are orthogonal with respect to the Jacobi weight function $w_{\alpha,\beta}(t) = (1-t)^\alpha (1+t)^\beta$ on $[-1,1]$, namely,

$$\int_{-1}^{1} P_n^{(\alpha,\beta)}(t)\, P_m^{(\alpha,\beta)}(t)\, w_{\alpha,\beta}(t)\, dt = \begin{cases} \dfrac{2^{\alpha+\beta+1}\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{(2n+\alpha+\beta+1)\Gamma(n+1)\Gamma(n+\alpha+\beta+1)}, & n=m, \\ 0, & n \neq m. \end{cases}$$

Here, as usual, Γ is the gamma function.

For $\lambda > -\frac{1}{2}$, $\mu \geq 0$, and $n \in \mathbb{N}_0$, the generalized Gegenbauer polynomials $C_n^{(\lambda,\mu)}(\cdot)$ are defined by

$$C_{2n}^{(\lambda,\mu)}(t) = a_{2n}^{(\lambda,\mu)} P_n^{(\lambda-1/2,\mu-1/2)}(2t^2-1), \qquad a_{2n}^{(\lambda,\mu)} = \frac{(\lambda+\mu)_n}{(\mu+\frac{1}{2})_n},$$

$$C_{2n+1}^{(\lambda,\mu)}(t) = a_{2n+1}^{(\lambda,\mu)}\, t P_n^{(\lambda-1/2,\mu+1/2)}(2t^2-1), \qquad a_{2n+1}^{(\lambda,\mu)} = \frac{(\lambda+\mu)_{n+1}}{(\mu+\frac{1}{2})_{n+1}},$$

where $(\lambda)_n$ denotes the Pochhammer symbol given by

$$(\lambda)_0 = 1, \quad (\lambda)_n = \lambda(\lambda+1)\cdots(\lambda+n-1) \quad \text{for} \quad n = 1, 2, \ldots.$$

They are orthogonal with respect to the weight function

$$v_{\lambda,\mu}(t) = |t|^{2\mu}(1-t^2)^{\lambda-1/2}, \quad t \in [-1,1]. \tag{1}$$

For $\mu = 0$, these polynomials, denoted by $C_n^\lambda(\cdot)$, are called the Gegenbauer polynomials:

$$C_n^\lambda(t) = C_n^{(\lambda,0)}(t) = \frac{(2\lambda)_n}{(\lambda+\frac{1}{2})_n} P_n^{(\lambda-1/2,\lambda-1/2)}(t).$$

For $\lambda > -\frac{1}{2}$, $\mu > 0$, and $n \in \mathbb{N}_0$, we have the following connection:

$$C_n^{(\lambda,\mu)}(t) = c_\mu \int_{-1}^{1} C_n^{\lambda+\mu}(tx)(1+x)(1-x^2)^{\mu-1}\,dx, \quad c_\mu^{-1} = 2\int_0^1 (1-x^2)^{\mu-1}\,dx.$$

Denote by $\bigl\{\widetilde{C}_n^{(\lambda,\mu)}(\cdot)\bigr\}_{n=0}^{\infty}$ the sequence of orthonormal generalized Gegenbauer polynomials. It is easily verified that these polynomials are given by the following formulae:

$$\begin{aligned}
&\widetilde{C}_{2n}^{(\lambda,\mu)}(t) = \widetilde{a}_{2n}^{(\lambda,\mu)} P_n^{(\lambda-1/2,\mu-1/2)}(2t^2-1),\\
&\widetilde{a}_{2n}^{(\lambda,\mu)} = \left(\frac{(2n+\lambda+\mu)\Gamma(n+1)\Gamma(n+\lambda+\mu)}{\Gamma(n+\lambda+\frac{1}{2})\Gamma(n+\mu+\frac{1}{2})}\right)^{1/2},\\
&\widetilde{C}_{2n+1}^{(\lambda,\mu)}(t) = \widetilde{a}_{2n+1}^{(\lambda,\mu)}\, t P_n^{(\lambda-1/2,\mu+1/2)}(2t^2-1),\\
&\widetilde{a}_{2n+1}^{(\lambda,\mu)} = \left(\frac{(2n+\lambda+\mu+1)\Gamma(n+1)\Gamma(n+\lambda+\mu+1)}{\Gamma(n+\lambda+\frac{1}{2})\Gamma(n+\mu+\frac{3}{2})}\right)^{1/2}.
\end{aligned}$$

The generalized Gegenbauer polynomials play an important role in Dunkl harmonic analysis (see, for example, [3, 4]). So, the study of these polynomials and their applications is very natural.

The notation $f(n) \asymp g(n)$, $n \to \infty$, means that there exist positive constants C_1, C_2, and a positive integer n_0 such that $0 \le C_1 g(n) \le f(n) \le C_2 g(n)$ for all $n \ge n_0$. For brevity, we will omit "$n \to \infty$" in the asymptotic notation.

Define the uniform norm of a continuous function f on $[-1,1]$ by

$$\|f\|_\infty = \max_{-1\le t\le 1} |f(t)|.$$

The maximum of two real numbers x and y is denoted by $\max(x,y)$.

In [8], we prove the following result.

Theorem 1. *Let $\lambda > -\frac{1}{2}$, $\mu > 0$. Then*

$$\bigl\|\widetilde{C}_n^{(\lambda,\mu)}\bigr\|_\infty \asymp n^{\max(\lambda,\mu)}.$$

Given $1 \le p \le \infty$, we denote by $L_p(v_{\lambda,\mu})$ the space of complex-valued Lebesgue measurable functions f on $[-1,1]$ with finite norm

$$\begin{aligned}
&\|f\|_{L_p(v_{\lambda,\mu})} = \Bigl(\textstyle\int_{-1}^{1} |f(t)|^p\, v_{\lambda,\mu}(t)\,dt\Bigr)^{1/p}, && 1 \le p < \infty,\\
&\|f\|_{L_\infty} = \operatorname*{ess\,sup}_{x\in[-1,1]} |f(x)|, && p = \infty.
\end{aligned}$$

For a function $f \in L_p(v_{\lambda,\mu})$, $1 \le p \le \infty$, the generalized Gegenbauer expansion is defined by

$$f(t) \sim \sum_{n=0}^{\infty} \hat{f}_n \widetilde{C}_n^{(\lambda,\mu)}(t), \qquad \text{where} \quad \hat{f}_n = \int_{-1}^{1} f(t)\,\widetilde{C}_n^{(\lambda,\mu)}(t)\,v_{\lambda,\mu}(t)\,dt.$$

For $1 < p < \infty$, we denote by p' the conjugate exponent to p, that is, $\frac{1}{p} + \frac{1}{p'} = 1$.

The aim of this paper is to establish Paley-type and Hausdorff – Young – Paley-type inequalities for generalized Gegenbauer expansions in Sections 2 and 3, respectively.

2. Paley-type inequalities for generalized Gegenbauer expansions

Using well-known techniques and the following obvious lemma, we establish in Theorem 2 Paley-type inequalities for generalized Gegenbauer expansions.

Lemma 1. *Suppose $A > 0$, $\gamma \geq 1$, φ is a positive function on $\mathbb{N}_0$, ψ is a non-negative function on $\mathbb{N}_0$, and*

$$\sup_{t>0}\Big\{t^{\gamma-1}\Big(\sum_{t\leq\varphi(n)\leq A}\psi(n)\Big)\Big\} < \infty.$$

Then

$$\sum_{\varphi(n)\leq A}(\varphi(n))^{\gamma}\psi(n) = \int_0^{A^\gamma}\Big(\sum_{\tau\leq(\varphi(n))^\gamma\leq A^\gamma}\psi(n)\Big)\,d\tau = \gamma\int_0^{A}t^{\gamma-1}\Big(\sum_{t\leq\varphi(n)\leq A}\psi(n)\Big)\,dt.$$

Theorem 2. *(a) If $1 < p \leq 2$, $f \in L_p(v_{\lambda,\mu})$, ω is a positive function on $\mathbb{N}_0$ such that*

$$M_\omega = \sup_{t>0}\Big\{t\Big(\sum_{\omega(n)\geq t}(n+1)^{2\max(\lambda,\mu)}\Big)\Big\} < \infty, \tag{2}$$

then

$$\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{p}-\frac{1}{p'}\right)\max(\lambda,\mu)}\,(\omega(n))^{\frac{1}{p}-\frac{1}{p'}}|\hat{f}_n|\Big)^p\Big\}^{1/p} \leq A_p M_\omega^{\frac{1}{p}-\frac{1}{p'}}\|f\|_{L_p(v_{\lambda,\mu})}. \tag{3}$$

(b) If $2 \leq q < \infty$, ω is a positive function on $\mathbb{N}_0$ satisfying (2) and ϕ is a non-negative function on $\mathbb{N}_0$ such that

$$\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{q}-\frac{1}{q'}\right)\max(\lambda,\mu)}(\omega(n))^{\frac{1}{q}-\frac{1}{q'}}|\phi(n)|\Big)^q < \infty, \tag{4}$$

then the algebraic polynomials

$$\Phi_N(t) = \sum_{n=0}^{N}\phi(n)\widetilde{C}_n^{(\lambda,\mu)}(t)$$

converge in $L_q(v_{\lambda,\mu})$ to a function f satisfying $\hat{f}_n = \phi(n)$, $n \in \mathbb{N}_0$, and

$$\|f\|_{L_q(v_{\lambda,\mu})} \leq A_{q'}M_\omega^{\frac{1}{q'}-\frac{1}{q}}\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{q}-\frac{1}{q'}\right)\max(\lambda,\mu)}(\omega(n))^{\frac{1}{q}-\frac{1}{q'}}|\phi(n)|\Big)^q\Big\}^{1/q}. \tag{5}$$

P r o o f. Let $\sigma = \max(\lambda,\mu)$.

(a) To prove (3), we note that for $p = 2$ the Parseval identity implies equality in (3) with $A_2 = 1$. We now consider (3) as the transformation from $L_p(v_{\lambda,\mu})$ into the sequence $\big\{(n+1)^{-\sigma}(\omega(n))^{-1}\hat{f}_n\big\}_{n=0}^{\infty}$ in the ℓ_p norm with the weight $\big\{(n+1)^{2\sigma}(\omega(n))^2\big\}_{n=0}^{\infty}$ and show

that this transformation is of weak type $(1,1)$. By Theorem 1, Lemma 1 and (2), we have $|\hat{f}_n| \le C\|f\|_{L_1(v_{\lambda,\mu})}(n+1)^{\sigma}$ and

$$m\{n\colon\ (n+1)^{-\sigma}(\omega(n))^{-1}|\hat{f}_n| > t\} \le m\{n\colon\ C(\omega(n))^{-1}\|f\|_{L_1(v_{\lambda,\mu})} > t\} \le$$
$$\le \sum_{\omega(n)\le A} (n+1)^{2\sigma}(\omega(n))^2 = 2\int_0^A t\Big(\sum_{t\le\omega(n)\le A}(n+1)^{2\sigma}\Big)\,dt \le 2AM_{\omega},$$

where $A = C\frac{\|f\|_{L_1(v_{\lambda,\mu})}}{t}$.

The last estimate is a weak $(1,1)$ estimate which, using the Marcinkiewicz interpolation theorem, implies (3).

(b) We have $1<q'\le 2$. For brevity, write ψ_n in place of $\Big((n+1)^{\left(\frac{1}{q}-\frac{1}{q'}\right)\sigma}(\omega(n))^{\frac{1}{q}-\frac{1}{q'}}|\phi(n)|\Big)^q$. Suppose that $g \in L_{q'}(v_{\lambda,\mu})$ and that $N < N'$ are positive integers. Applying Hölder's inequality and (a), we find that

$$\begin{aligned}
\Big|\int_{-1}^{1}\Phi_N(t)\,g(t)\,v_{\lambda,\mu}(t)\,dt\Big| &= \Big|\sum_{n=0}^{N}\phi(n)\hat{g}_n\Big| = \\
&= \Big|\sum_{n=0}^{N}\Big((n+1)^{\left(\frac{1}{q}-\frac{1}{q'}\right)\sigma}(\omega(n))^{\frac{1}{q}-\frac{1}{q'}}\phi(n)\Big)\Big((n+1)^{\left(\frac{1}{q'}-\frac{1}{q}\right)\sigma}(\omega(n))^{\frac{1}{q'}-\frac{1}{q}}\hat{g}_n\Big)\Big| \le \\
&\le \Big\{\sum_{n=0}^{N}\psi_n\Big\}^{1/q}\Big\{\sum_{n=0}^{N}\Big((n+1)^{\left(\frac{1}{q'}-\frac{1}{q}\right)\sigma}(\omega(n))^{\frac{1}{q'}-\frac{1}{q}}|\hat{g}_n|\Big)^{q'}\Big\}^{1/q'} \le \\
&\le \Big\{\sum_{n=0}^{N}\psi_n\Big\}^{1/q} A_{q'}M_{\omega}^{\frac{1}{q'}-\frac{1}{q}}\|g\|_{L_{q'}(v_{\lambda,\mu})}.
\end{aligned} \tag{6}$$

Similarly,

$$\Big|\int_{-1}^{1}\big(\Phi_N(t)-\Phi_{N'}(t)\big)\,g(t)\,v_{\lambda,\mu}(t)\,dt\Big| \le \Big\{\sum_{n=N+1}^{N'}\psi_n\Big\}^{1/q} A_{q'}M_{\omega}^{\frac{1}{q'}-\frac{1}{q}}\|g\|_{L_{q'}(v_{\lambda,\mu})}. \tag{7}$$

Hence, by [5, Theorem (12.13)], the inequalities (6) and (7) lead respectively to the estimates

$$\|\Phi_N\|_{L_q(v_{\lambda,\mu})} \le \Big\{\sum_{n=0}^{N}\psi_n\Big\}^{1/q} A_{q'}M_{\omega}^{\frac{1}{q'}-\frac{1}{q}} \tag{8}$$

and

$$\|\Phi_N-\Phi_{N'}\|_{L_q(v_{\lambda,\mu})} \le \Big\{\sum_{n=N+1}^{N'}\psi_n\Big\}^{1/q} A_{q'}M_{\omega}^{\frac{1}{q'}-\frac{1}{q}}.$$

The last inequality combined with (4) shows that the sequence $\{\Phi_N\}_{N=1}^{\infty}$ is a Cauchy sequence in $L_q(v_{\lambda,\mu})$ and therefore convergent in $L_q(v_{\lambda,\mu})$; let f be its limit. Then, by mean convergence,

$$\hat{f}_n = \lim_{N\to\infty}\big(\widehat{\Phi_N}\big)_n, \quad n\in\mathbb{N}_0,$$

which is easily seen to equal $\phi(n)$. Moreover, the defining relation

$$f = \lim_{N\to\infty} \Phi_N \qquad \text{in} \quad L_q(v_{\lambda,\mu})$$

and the inequality (8) show that (5) holds and so complete the proof. □

3. Hausdorff – Young – Paley-type inequalities for generalized Gegenbauer expansions

In [9], we prove the following Hausdorff – Young-type inequalities for generalized Gegenbauer expansions.

Theorem 3. (*a*) *If* $1 < p \le 2$ *and* $f \in L_p(v_{\lambda,\mu})$, *then*

$$\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{p'}-\frac{1}{p}\right)\max(\lambda,\mu)}|\hat{f}_n|\Big)^{p'}\Big\}^{1/p'} \le B_p\,\|f\|_{L_p(v_{\lambda,\mu})}.$$

(*b*) *If* $2 \le q < \infty$ *and* ϕ *is a function on non-negative integers satisfying*

$$\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{q'}-\frac{1}{q}\right)\max(\lambda,\mu)}|\phi(n)|\Big)^{q'} < \infty,$$

then the algebraic polynomials

$$\Phi_N(t) = \sum_{n=0}^{N} \phi(n)\,\widetilde{C}_n^{(\lambda,\mu)}(t)$$

converge in $L_q(v_{\lambda,\mu})$ *to a function* f *satisfying* $\hat{f}_n = \phi(n)$, $n \in \mathbb{N}_0$, *and*

$$\|f\|_{L_q(v_{\lambda,\mu})} \le B_{q'}\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{q'}-\frac{1}{q}\right)\max(\lambda,\mu)}|\phi(n)|\Big)^{q'}\Big\}^{1/q'}.$$

Theorem 5 contains the Paley-type and the Hausdorff – Young-type inequalities for the expansions by orthonormal polynomials with respect to the weight function $v_{\lambda,\mu}$ (see (1)). To prove it, we need Stein's modification of the Riesz – Thorin interpolation theorem (see [6, Theorem 2, p. 485]) given below.

Theorem 4 (Stein). *Suppose* ν_1 *and* ν_2 *are* σ*-finite measures on* M *and* S, *respectively, and* T *is a linear operator defined on* ν_1*-measurable functions on* M *to* ν_2*-measurable functions on* S. *Let* $1 \le r_0, r_1, s_0, s_1 \le \infty$ *and* $\frac{1}{r} = \frac{1-t}{r_0} + \frac{t}{r_1}$, $\frac{1}{s} = \frac{1-t}{s_0} + \frac{t}{s_1}$, *where* $0 \le t \le 1$. *Suppose further that*

$$\|(Tg)\cdot v_i\|_{L_{s_i}(S,\nu_2)} \le L_i\|g\cdot u_i\|_{L_{r_i}(M,\nu_1)}, \quad i = 0, 1,$$

where u_i *and* v_i *are non-negative weight functions. Let* $u = u_0^{1-t}\cdot u_1^t$, $v = v_0^{1-t}\cdot v_1^t$.

Then

$$\|(Tg)\cdot v\|_{L_s(S,\nu_2)} \le L\|g\cdot u\|_{L_r(M,\nu_1)}$$

with $L = L_0^{1-t}\cdot L_1^t$.

Theorem 5. *Let* $\sigma = \max(\lambda, \mu)$.

(*a*) *If* $1 < p \le 2$, $f \in L_p(v_{\lambda,\mu})$, ω *is a positive function on* $\mathbb{N}_0$ *satisfying the condition* (2), *and* $p \le s \le p'$, *then*

$$\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{(\frac{2}{s}-1)\sigma}(\omega(n))^{\frac{1}{s}-\frac{1}{p'}}|\hat{f}_n|\Big)^s\Big\}^{1/s} \le C_p(s)\, M_\omega^{\frac{1}{s}-\frac{1}{p'}}\, \|f\|_{L_p(v_{\lambda,\mu})}. \tag{9}$$

(*b*) *If* $2 \le q < \infty$, $q' \le r \le q$, ω *is a positive function on* $\mathbb{N}_0$ *satisfying the condition* (2) *and* ϕ *is a non-negative function on* $\mathbb{N}_0$ *such that*

$$\sum_{n=0}^{\infty}\Big((n+1)^{(1-\frac{2}{r})\sigma}(\omega(n))^{\frac{1}{q}-\frac{1}{r}}|\phi(n)|\Big)^{r'} < \infty,$$

then the algebraic polynomials

$$\Phi_N(t) = \sum_{n=0}^{N}\phi(n)\widetilde{C}_n^{(\lambda,\mu)}(t)$$

converge in $L_q(v_{\lambda,\mu})$ *to a function* f *satisfying* $\hat{f}_n = \phi(n)$, $n \in \mathbb{N}_0$, *and*

$$\|f\|_{L_q(v_{\lambda,\mu})} \le C_{q'}(r) M_\omega^{\frac{1}{r}-\frac{1}{q}}\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{(1-\frac{2}{r})\sigma}(\omega(n))^{\frac{1}{q}-\frac{1}{r}}|\phi(n)|\Big)^{r'}\Big\}^{1/r'}.$$

P r o o f. (a) This part was proved for $s = p$ (with $C_p(p) = A_p$) and $s = p'$ (with $C_p(p') = B_p$) in Theorems 2 and 3, respectively. So for $p = 2$, we obtain the equality in (9) with $C_2(2) = 1$.

Consider now the case that $1 < p < 2$. To prove (9), we set in Theorem 4: $M = [-1, 1]$, ν_1 the Lebesgue measure, $S = \{n\}_{n=0}^{\infty}$, ν_2 the counting measure, $g = f$, $Tg = \{\hat{f}_n\}_{n=0}^{\infty}$, $r = r_0 = r_1 = p$, $u = u_0 = u_1 = v_{\lambda,\mu}$, $s_0 = p'$, $s_1 = p$, $v_0 = \big\{(n+1)^{\left(\frac{1}{p'}-\frac{1}{p}\right)\sigma}\big\}_{n=0}^{\infty}$, $v_1 = \big\{(n+1)^{\left(\frac{1}{p}-\frac{1}{p'}\right)\sigma}(\omega(n))^{\frac{1}{p}-\frac{1}{p'}}\big\}_{n=0}^{\infty}$, $L_0 = B_p$, $L_1 = A_p M_\omega^{\frac{1}{p}-\frac{1}{p'}}$, and $\frac{1}{s} = \frac{1-t}{p'} + \frac{t}{p}$. As $\frac{1}{p} + \frac{1}{p'} = 1$, $\frac{1}{s} - \frac{1}{p} = (1-t)\big(\frac{1}{p'} - \frac{1}{p}\big)$, $\frac{1}{p'} - \frac{1}{s} = t\big(\frac{1}{p'} - \frac{1}{p}\big)$, the proof of (9) is concluded.

Because of

$$1 - t = \frac{\frac{1}{s} - \frac{1}{p}}{\frac{1}{p'} - \frac{1}{p}}, \quad t = \frac{\frac{1}{p'} - \frac{1}{s}}{\frac{1}{p'} - \frac{1}{p}},$$

it is clear that $C_p(s) = B_p^{1-t} A_p^t$ and $M_\omega^{\left(\frac{1}{p}-\frac{1}{p'}\right)t} = M_\omega^{\frac{1}{s}-\frac{1}{p'}}$.

(b) Taking into account the previously given proof (see part (b) in Theorem 2), the proof is obvious and left to the reader. □

4. Conclusion

As an application of the results above, we are going to obtain sufficient condition for a special sequence of positive real numbers to be a Fourier multiplier.

Acknowledgements

The author would like to thank Michael Ruzhansky for the idea of this work (see the e-print [1]).

REFERENCES

1. R. Akylzhanov, E. Nursultanov, and M. Ruzhansky, Hardy – Littlewood, Hausdorff – Young – Paley inequalities, and L^p-L^q Fourier multipliers on compact homogeneous manifolds, arXiv preprint 1504.07043 (2015).
2. G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications **71**, Cambridge University Press, Cambridge, 1999.
3. F. Dai and Y. Xu, *Approximation theory and harmonic analysis on spheres and balls*, Springer Monographs in Mathematics, Springer, 2013.
4. C. F. Dunkl and Y. Xu, *Orthogonal polynomials of several variables*, 2nd ed., Encyclopedia of Mathematics and its Applications **155**, Cambridge University Press, Cambridge, 2014.
5. E. Hewitt and K. A. Ross, *Abstract harmonic analysis. Vol. I*, Springer-Verlag, Heidelberg, 1963.
6. E. Stein, Interpolation of linear operators, *Trans. Amer. Math. Soc.* **83** (1956), 482–492.
7. G. Szegö, *Orthogonal polynomials*, 4th ed., American Mathematical Society Colloquium Publications **23**, American Mathematical Society, Providence, Rhode Island, 1975.
8. R. A. Veprintsev, On the asymptotic behavior of the maximum absolute value of generalized Gegenbauer polynomials, arXiv preprint 1602.01023 (2016).
9. R. A. Veprintsev, On Hardy – Littlewood-type and Hausdorff – Young-type inequalities for generalized Gegenbauer expansions, arXiv preprint 1602.04326 (2016).

Independent researcher, Uzlovaya, Russia
E-mail address: **veprintsevroma@gmail.com**